

**WHAT IS CLAIMED IS:**

1. A method of identifying a candidate substance that inhibits the aggregation of an aggregate-prone amyloid protein, comprising:

- 5
- Sub B2
- (a) contacting a yeast cell that expresses an aggregate-prone amyloid protein with said candidate substance under conditions effective to allow aggregated amyloid formation; and
- 10
- (b) determining the ability of said candidate substance to inhibit the aggregation of the aggregate-prone amyloid protein.

15

2. The method of claim 1, wherein the aggregate-prone amyloid protein comprises a Sup35 or URE3 polypeptide.

20

3. The method of claim 1, wherein the aggregate-prone amyloid protein comprises a PrP or  $\beta$ -amyloid polypeptide.

25

4. The method of claim 1, wherein the aggregate-prone amyloid protein is a chimeric protein.

5. The method of claim 4, wherein the chimeric protein comprises at least the N-terminal domain of Sup35.

30

6. The method of claim 4, wherein the chimeric protein comprises at least an aggregate forming domain of a mammalian amyloid polypeptide.

Sup B4  
5 7. The method of claim 4, wherein the chimeric protein comprises at least an aggregate forming domain of an aggregate-prone amyloid protein operably attached to a detectable marker protein.

8. The method of claim 7, wherein said marker protein is green fluorescent protein or luciferase.

10

9. The method of claim 7, wherein said marker protein is a drug-resistance marker protein.

10. The method of claim 7, wherein said marker protein is a hormone receptor.

11. The method of claim 10, wherein said hormone receptor is a glucocorticoid receptor.

Sup B5  
12. The method of claim 6, wherein the mammalian amyloid polypeptide is PrP or  $\beta$ -amyloid.

25 13. The method of claim 12, wherein the chimeric protein comprises as least about amino acids 1-42 of  $\beta$ -amyloid protein.

Sup E4  
30 14. The method of claim 4, wherein the chimeric protein comprises Sup35 in which the N-terminal domain has been replaced by amino acids 1-42 of  $\beta$ -amyloid protein.

15. The method of claim 1, wherein any aggregation of the aggregate-prone amyloid protein is detected by the ability of the aggregated protein to bind Congo Red.

16. The method of claim 1, wherein any aggregation of the aggregate-prone amyloid protein is detected by increased protease resistance of the aggregated protein.

17. The method of claim 1, wherein the aggregate-prone amyloid protein is labeled.

18. The method of claim 17, wherein the label is a radioactive isotope, a fluorophore, or a chromophore.

19. The method of claim 18, wherein the label is  $^{35}\text{S}$ .

20. The method of claim 18, wherein the fluorophore comprises a green fluorescent protein polypeptide.

21. The method of claim 1, wherein any aggregation is determined by the presence of a [PSI<sup>+</sup>] phenotype.

22. The method of claim 1, wherein said yeast cell overexpresses Hsp104.

23. A method of identifying a candidate substance for therapeutic activity against an amyloidogenic disease in an animal, said method comprising:

(a) contacting a yeast cell that expresses an aggregate-prone amyloid protein with said candidate substance under conditions effective to allow amyloid formation; and

(b) determining the ability of said candidate substance to inhibit aggregation of the aggregate-prone amyloid protein,

wherein the ability to inhibit aggregation is indicative of therapeutic activity.

24. The method of claim 23, wherein the aggregate-prone amyloid protein comprises a PrP,  $\beta$ -amyloid, Sup35, or URE3 polypeptide.

25. The method of claim 23, wherein the protein is a chimeric protein.

26. The method of claim 25, wherein the chimeric protein comprises a Sup35 polypeptide.

27. The method of claim 25, wherein the chimeric protein comprises a mammalian amyloid polypeptide.

28. The method of claim 27, wherein the mammalian amyloid polypeptide is PrP or  $\beta$ -amyloid.

29. The method of claim 23, wherein any aggregation of the aggregate-prone amyloid protein is detected by the ability of the aggregation to bind Congo Red.

30. The method of claim 23, wherein the aggregate-prone amyloid protein is labeled.

31. The method of claim 30, wherein the label is a radioactive isotope, a fluorophore, or a chromophore.

32. The method of claim 31, wherein the label is <sup>35</sup>S.

33. The method of claim 31, wherein the fluorophore comprises a green fluorescent protein polypeptide.

34. The method of claim 23, wherein the aggregation of the aggregate-prone amyloid protein is determined by the presence of a [PSI<sup>+</sup>] phenotype.

35. The method of claim 23, wherein the disease is selected from the group consisting of Alzheimer's disease, scrapie, spongiform encephalopathy in a mammal, kuru, Creutzfeldt-Jakob disease, Gestmann-Strausser-Scheinker disease, or fatal familial insomnia.

36. The method of claim 35, wherein the mammal is bovine, feline, a mink, deer, elk, a mouse, a hamster, an ape, a monkey, or human.

Ad B7

Add D3